



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

NOTES ON THE TERTIARY GEOLOGY AND PALEONTOLOGY OF THE
SOUTHERN UNITED STATES.

BY PROF. ANGELO HEILPRIN.

Eocene of Texas.—In a limited collection of fossils from near the northern border of San Augustine Co., Texas, transmitted to me for examination by the Texas State Geological and Scientific Association, I have been able to determine the following species:—

<i>Ostrea Alabamiensis.</i>	<i>Cardita Blandingi (alticosta).</i>
<i>Ostrea sellaformis.</i>	<i>Crassatella antestriata.</i>
<i>Ostrea divaricata.</i>	<i>Corbula Texana.</i>
<i>Pecten Deshayesii.</i>	<i>Buccitriton altile? (Texana).</i>
<i>Anomia?</i>	<i>Scalaria</i> sp. indet.

The horizon represented is evidently the "Claibornian," the deposits probably occupying a position in the "Jacksonian" area.

The Nummulitic of Florida.—Prof. G. A. Wetherby, of Cincinnati, has furnished me with a number of rock fragments obtained at a locality some six miles southwest of Gainesville, Fla. They are interesting as containing, in addition to one or more species of *Orbitoides*, several nearly perfect individuals of *Nummulites Floridanus* Heilprin, which, therefore, represent the most northern locality in the State where the members of this group of Foraminifera have thus far been found. They lend further confirmation to the views already advanced by the writer as to the broad extent of the southern Nummulitic formation, and to the relative antiquity of the Floridan peninsula. One considerable fragment of a *Heterostegina* is also represented in the rock.

Since the receipt of Prof. Wetherby's specimens I have been favored, through Mr. Joseph Willcox, of this city, with other fragments from approximately the same locality, Arredonda, Alachua Co., which also contain *Nummulites Floridanus*, *Orbitoides*, and *Operculina rotella* (*O. complanata?*).

Eocene of Kentucky.—Mr. R. H. Loughridge has kindly forwarded to me for determination a number of fossils collected by the Geological Survey of Kentucky, from the immediate neighborhood of Paducah. They are mainly in the form of casts in a highly ferruginous and fairly micaceous yellow-white sandstone,